

Estimation of the volume of an excursion set of a Gaussian process using intrinsic Kriging

Emmanuel Vazquez^a

E-mail address: emmanuel.vazquez@supelec.fr

Miguel Piera Martínez^{a,b}

a: Département Signaux et Systèmes Électroniques, Supélec, 91192 Gif-sur-Yvette, France

b: Laboratoire des Signaux et Systèmes, CNRS, Supélec, Université Paris-Sud, 91192 Gif-sur-Yvette, France

February 2, 2008

Abstract — Assume that a Gaussian process ξ is predicted from n pointwise observations by intrinsic Kriging and that the volume of the excursion set of ξ above a given threshold u is approximated by the volume of the predictor. The first part of this paper gives a bound on the convergence rate of the approximated volume. The second part describes an algorithm that constructs a sequence of points to yield a fast convergence of the approximation. The estimation of the volume of an excursion set is a highly relevant problem for the industrial world since it corresponds to the estimation of the failure probability of a system that is known only through sampled observations.

Keywords – Excursion set; Gaussian process; Intrinsic Kriging; Quantile estimation; Failure probability; Design of experiments

1 Introduction

The problem to be considered in this paper is the estimation of the probability

$$\mathcal{P}_u := \mathbf{P}\{f(X) \geq u\}, \quad (1)$$

where $f(x)$ is a real function defined over an arbitrary set $\mathbb{X}$ ($\mathbb{X} = [0, 1]^d$ or $\mathbb{X} = \mathbb{R}^d$, in most situations) endowed with a probability measure μ and $X \in \mathbb{X}$ is a random vector with the distribution μ . In practice, the estimation of (1) is based on a finite sequence of evaluations of f at points $(x_i)_{1 \leq i \leq n}$ in $\mathbb{X}$. Another way of looking at (1) is via the excursion set

$$A_u(f) := \{x \in \mathbb{X} : f(x) \geq u\} \quad (2)$$

of the function f above the level u , since $\mathbf{P}\{f(X) \geq u\}$ is the volume $\mu(A_u(f))$, hereafter denoted by $|A_u(f)|$.

Such a problem is frequently encountered in engineering: the probability that the inputs of the system will generate a level of a function of the outputs that exceeds a specified reference level may be expressed as (1) (where in this case, X is the vector of the inputs of the system and f is a statistic of the outputs). Since to obtain the value of f at a given x may be very expensive in practice, because it may involve heavy computer codes for instance, it is often essential to estimate $\mathcal{P}_u$ using as few evaluations of f as possible.

To overcome the problem of evaluating f many times, one possible approach is to estimate $|A_u(f_n)|$ instead of $|A_u(f)|$, where f_n is an approximation of f constructed from a small set $\{f(x_1), \dots, f(x_n)\}$ of pointwise evaluations. Such an approximation can be obtained by assuming that f is a sample path of a Gaussian random process ξ and by using a linear predictor ξ_n of ξ constructed from $\xi(x_i)$, $i = 1, \dots, n$. In this paper, intrinsic Kriging (Matheron, 1973) will be used to obtain ξ_n . We shall show in Section 2 that this method is likely to give faster convergences than the classical Monte Carlo estimators, depending on the regularity of ξ .

A second step is to choose a sequence of evaluation points (x_i) so that $|A_u(\xi_n)| - |A_u(\xi)|$ conditioned on the random variables $\xi(x_i)$, $i \leq n$, converges rapidly to zero. Section 3 presents an acceleration algorithm based on computing an upper bound of the mean square error of volume approximation conditioned on the events $\{\xi(x_i) = f(x_i), i = 1 \dots, n\}$: a point x_{n+1} is selected so that evaluating $f(x_{n+1})$ yields the potential largest decrease of the upper bound. Section 4 provides a numerical example.

2 Excursion set volume estimation by intrinsic Kriging

This section deals with the estimation of the probability $\mathcal{P}_u$ from observations of f at a finite sequence of points $(x_i)_{1 \leq i \leq n}$. As mentioned above, $\mathcal{P}_u$ is the volume of $A_u(f)$ under the probability distribution μ . We assume moreover that f is a sample path of a (separable) Gaussian process ξ , with mean $m(x)$, $x \in \mathbb{X}$, and covariance $k(x, y)$, $(x, y) \in \mathbb{X}^2$.

2.1 Monte Carlo estimation

Monte Carlo is a commonly used method to estimate $|A_u(\xi)|$. The volume of excursion of a Gaussian process ξ may be estimated by

$$|A_u(\xi)|_l := \frac{1}{l} \sum_{i=1}^l \mathbf{1}_{\{\xi(X_i) \geq u\}} \xrightarrow{l} |A_u(\xi)| \quad \text{a.s.} \quad (3)$$

where the X_i s are independent random variables with distribution μ . The estimator (3) is unbiased, since $E[|A_u(\xi)|_l | \xi] = |A_u(\xi)|$, and

$$E[(|A_u(\xi)|_l - |A_u(\xi)|)^2 | \xi] = \frac{1}{l} |A_u(\xi)| (1 - |A_u(\xi)|) .$$

If evaluating f (a sample path of ξ) at many points of $\mathbb{X}$ is not particularly demanding, then estimating $|A_u(f)|$ is straightforward. However, if $|A_u(f)|$ is small, then the variance of the Monte Carlo estimator is approximately

$|A_u(f)|/l$. To achieve a given standard deviation $\kappa|A_u(f)|$, with $\kappa > 0$ small, the required number of evaluations is approximately $1/(\kappa^2|A_u(f)|)$, i.e. it is high. Thus, the convergence of (3) may be too slow in many real applications where doing a lot of evaluations of f may not be affordable (for instance, f may be a complex computer simulation and may take hours or days to run). Of course, many other methods have been proposed to improve the basic Monte Carlo convergence. For instance, methods based on importance sampling, on cross-entropy (Rubinstein, 1999), on the classical extreme value theory (e.g. Embrechts et al., 1997), etc. They are not considered here for the sake of brevity.

2.2 Estimation based on an approximation

An alternative approach is to replace f by an approximation f_n constructed from a set of n point evaluations of f . Provided f_n converges rapidly enough to f , one expects a good estimation of the excursion sets and their volume using only a few evaluations of f . There are many ways of constructing such an approximation. Let us mention two classical methods: regularized regressions in reproducing kernel Hilbert spaces, e.g. splines or radial basis functions (see for instance Wendland, 2005), and linear prediction of random processes, also known as Kriging (see for instance Chilès and Delfiner, 1999). In this paper, we shall adopt the probabilistic framework¹.

Thus, let us consider that an unbiased linear estimator ξ_n of ξ has been obtained from $\xi(x_1), \dots, \xi(x_n)$. In particular, we can use ordinary Kriging when the mean of $\xi(x)$ is known and intrinsic Kriging when it is unknown, which is more often the case.

Can we expect a faster convergence when ξ is replaced by ξ_n ? Here, we assume the computation time to evaluate $\xi_n(x)$, $x \in \mathbb{X}$, conditioned on $\xi(x_i) = f(x_i)$, $i = 1, \dots, n$, is small, which means that we can make $|A_u(\xi_n)|_l - |A_u(\xi_n)|$ negligible with respect to $|A_u(\xi_n)| - |A_u(\xi)|$. Thus, we are now interested in the convergence of $|A_u(\xi_n)|$ to $|A_u(\xi)|$. Section 2.2.2 shows how the convergence rate in mean square of $|A_u(\xi_n)|$ to $|A_u(\xi)|$ depends on the fill distance of $\mathbb{X}$ and the regularity of ξ . In Section 3, we shall propose an algorithm to speed up this rate by a sequential choice of the evaluation points.

2.2.1 Intrinsic Kriging basics

In this paper, we use *intrinsic Kriging* (IK) to obtain a linear predictor of ξ based on a finite set of pointwise observations of the process. We recall here the main results (Matheron, 1973). IK extends linear prediction when the mean of $\xi(x)$ is unknown but can be written as a linear parametric function $m(x) = b^\top p(x)$. Here, $p(x)$ is a q -dimensional vector of base functions of a vector space $\mathcal{N}$ of translation-stable functions (in practice, all polynomials of degree less or equal to l) and b is a vector of unknown parameters. Intrinsic Kriging assumes that observed values of f are samples from a representation of an intrinsic random function (IRF), a generalized random process defined over a space Λ_l of measures orthogonal to $\mathcal{N}$, and characterized by its stationary generalized covariance $k(h)$ (see the Appendix Section for more details).

¹In fact, these two classes of methods, which have been studied separately, are equivalent (see for instance Kimeldorf and Wahba (1970)).

Proposition 1 (Intrinsic Kriging, Matheron 1973). *Let ξ_G be an $IRF(l)$, with generalized covariance $k(h)$. Assume n observations be sample values of the random variables $\xi_{x_i}^{\text{obs}} = \xi(x_i) + N_i$, $i = 1, \dots, n$, where ξ is an unknown representation of ξ_G and the N_i s are zero-mean random variables independent of $\xi(x)$, with covariance matrix K_N .*

The intrinsic Kriging predictor of $\xi(x)$ based on the observations, is the linear projection $\xi_n(x) = \sum_i \lambda_{i,x} \xi_{x_i}^{\text{obs}}$ of $\xi(x)$ onto $\mathcal{H}_S = \text{span}\{\xi_{x_i}^{\text{obs}}, i = 1, \dots, n\}$, such that the variance of the prediction error $\xi(x) - \xi_n(x)$ is minimized under the constraint $\delta_x - \sum \lambda_{i,x} \delta_{x_i} \in \Lambda_l$. The coefficients $\lambda_{i,x}$, $i = 1, \dots, n$, are solutions of a system of linear equations, which can be written in matrix form as

$$\begin{pmatrix} K + K_N & P^\top \\ P & 0 \end{pmatrix} \begin{pmatrix} \lambda_x \\ \mu \end{pmatrix} = \begin{pmatrix} k_x \\ p_x \end{pmatrix}, \quad (4)$$

where K is the $n \times n$ matrix of generalized covariances $k(x_i - x_j)$, P is a $q \times n$ matrix with entries x_j^i for $j = 1, \dots, n$ and multi-indexes $i = (i_1, \dots, i_d)$ such that $|i| := i_1 + \dots + i_d \leq l$, μ is a vector of Lagrange coefficients, k_x is a vector of size n with entries $k(x - x_i)$ and p_x is a vector of size q with entries x^i , i such that $|i| \leq l$.

The variance of the prediction error is given by $\sigma_n(x)^2 := \text{var}[\xi(x) - \xi_n(x)] = k(0) - \lambda_x^\top k_x - \mu^\top p_x$.

Proof. See Matheron (1973). $\square$

2.2.2 Asymptotics

In this section, we shall justify that modeling the unknown f by a Gaussian random process ξ and estimating $|A_u(\xi)|$ by $|A_u(\xi_n)|$ is well-founded. Our objective is to establish a mean square convergence when the evaluation points fill $\mathbb{X}$.

Classical results in approximation theory (see for instance Wu and Schaback, 1993 ; Light and Wayne, 1998 ; Narcowich et al., 2003 ; Wendland, 2005) assert that the variance $\sigma_n^2(x)$ of the IK prediction error at x decreases as the sampling density or the regularity of the covariance increases. More precisely, if $\mathbb{X}$ is a bounded domain of $\mathbb{R}^d$, and the Fourier transform of $k(h)$, $h \in \mathbb{R}^d$, satisfies

$$c_1(1 + \|\omega\|_2^2)^{-\nu} \leq \tilde{k}(\omega) \leq c_2(1 + \|\omega\|_2^2)^{-\nu}.$$

with $\nu > d/2$, then

$$\|\sigma_n(\cdot)\|_\infty \leq C h_n^{\nu-d/2}, \quad (5)$$

where $h_n = \sup_{y \in \mathbb{X}} \min_i \|y - x_i\|_2$ is a *fill distance* of $(x_1, \dots, x_n)$ in $\mathbb{X}$.

The following theorem shows that a similar result holds for the process thresholded at a level u .

Theorem 1. *Let ξ be an unknown representation of an $IRF(l)$ ξ_G , and $\xi_n(x)$ be the IK predictor of ξ based on observations $\xi(x_i)$, $i = 1, \dots, n$. Define $\sigma_n(x) := \text{var}[\xi(x) - \xi_n(x)]^{1/2}$. Then,*

$$\mathbb{E} \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \right] = O(\sigma_n(x) |\log(\sigma_n(x))|^{1/2}) \quad \text{when } \sigma_n(x) \rightarrow 0.$$

Proof. For all $x \in \mathbb{X}$, $\xi(x) - \xi_n(x)$ is Gaussian with zero-mean and variance $\sigma_n(x)^2$ (but is not orthogonal to $\xi_n(x)$, as would be the case if the mean of ξ were known). Thus, $\forall x \in \mathbb{X}$ and $\forall n \in \mathbb{N}$, we can write $\xi(x)$ as

$$\xi(x) = (1 + a_n(x))\xi_n(x) + b_n(x) + \zeta_n(x), \quad (6)$$

where $a_n(x), b_n(x) \in \mathbb{R}$, $\zeta_n(x)$ is Gaussian and such that $E[\xi_n(x)\zeta_n(x)] = 0$ and $E[\zeta_n(x)] = 0$. This decomposition exists and is unique for every n . (To simplify notations, from now on, we shall omit the dependence on x when there is no ambiguity.)

Clearly, $\text{var}[\xi_n]$ is non-decreasing and can be assumed to be strictly positive for n large enough. Since $E[a_n\xi_n] = -b_n$, we have

$$\sigma_n^2 = \text{var}[a_n\xi_n + b_n + \zeta_n] = E[(a_n\xi_n + b_n + \zeta_n)^2] = a_n^2 \text{var}[\xi_n] + E[\zeta_n^2], \quad (7)$$

and thus, the following upper bounds hold for n large enough:

$$\begin{cases} |a_n| \leq K_a \sigma_n, & |b_n| \leq K_b \sigma_n, \\ \tilde{\sigma}_n := E[\zeta_n^2]^{1/2} \leq \sigma_n, \end{cases} \quad (8)$$

for some $K_a, K_b > 0$.

For some threshold $u \in \mathbb{R}$, let α be such that

$$\alpha > |a_n u + b_n| \geq 0, \quad (9)$$

and let $N \in \mathbb{N}$ be such that $\forall n > N$, $|a_n| < 1$. For all $n > N$, define

$$\begin{cases} h_n^- &= \frac{u - b_n - \alpha}{1 + a_n}, \\ h_n^+ &= \frac{u - b_n + \alpha}{1 + a_n}. \end{cases}$$

Note that $h_n^- < u < h_n^+$ and that

$$h_n^+ - h_n^- = \frac{2\alpha}{1 + a_n}.$$

For all $n > N$,

$$\begin{aligned} E \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \mid \xi_n(x) \right] &= \Psi \left(\frac{u - (1 + a_n)\xi_n - b_n}{\tilde{\sigma}_n} \right) \mathbf{1}_{\xi_n(x) < u} \\ &\quad + \Psi \left(-\frac{u - (1 + a_n)\xi_n - b_n}{\tilde{\sigma}_n} \right) \mathbf{1}_{\xi_n(x) \geq u}, \end{aligned} \quad (10)$$

in which Ψ denotes the tail of the standard Gaussian distribution function. Since

$$\begin{cases} \xi_n < h_n^- &\Rightarrow u - (1 + a_n)\xi_n - b_n > \alpha, \\ \xi_n > h_n^+ &\Rightarrow -u + (1 + a_n)\xi_n + b_n > \alpha, \end{cases}$$

and $\tilde{\sigma}_n \leq \sigma_n$, we have

$$E \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \mid \xi_n(x) \right] \leq \Psi \left(\frac{\alpha}{\sigma_n} \right) \mathbf{1}_{\xi_n(x) \in \mathbb{R} \setminus [h_n^-, h_n^+]} + \mathbf{1}_{\xi_n(x) \in [h_n^-, h_n^+]}. \quad (11)$$

By integrating with respect to the density of ξ_n , we obtain

$$\mathbb{E} \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \right] \leq \Psi \left(\frac{\alpha}{\sigma_n} \right) + c_0 \frac{2\alpha}{1 + a_n} \quad (12)$$

$$\leq \frac{\sigma_n}{\alpha \sqrt{2\pi}} \exp\left(-\frac{\alpha^2}{2\sigma_n^2}\right) + c_1 \alpha \quad (13)$$

where (13) uses a standard Gaussian tail inequality.

The upper bound can be tightened by replacing α with a sequence (α_n) such that

$$\alpha_n := \sqrt{2}\sigma_n |\log(\sigma_n)|^{1/2},$$

which satisfies (9) for n large enough. Therefore,

$$\mathbb{E} \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \right] \leq O(\sigma_n |\log(\sigma_n)|^{1/2}) \quad \text{when } \sigma_n \rightarrow 0. \quad (14)$$

□

Hence, if $\mathbb{X}$ is bounded:

$$\begin{aligned} \mathbb{E} \left[(|A_u(\xi)| - |A_u(\xi_n)|)^2 \right] &= \mathbb{E} \left[\left(\int_{\mathbb{X}} \mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u} d\mu \right)^2 \right] \\ &\leq \int_{\mathbb{X}} \mathbb{E} \left[(\mathbf{1}_{\xi(x) \geq u} - \mathbf{1}_{\xi_n(x) \geq u})^2 \right] d\mu \\ &\leq C \|\sigma_n(\cdot)\|_{\infty} |\log \|\sigma_n(\cdot)\|_{\infty}|^{1/2} \end{aligned} \quad (15)$$

when $n \rightarrow 0$ and $\|\sigma_n(\cdot)\|_{\infty} \rightarrow 0$.

Therefore, this simple result shows that the mean square convergence of $|A_u(\xi_n)|$ to $|A_u(\xi)|$ is related to the mean square convergence of ξ_n to ξ , hence, due to (5), to the regularity of the covariance and the fill-in distance of $\mathbb{X}$. Informally speaking, we can say that using an approximation will be more efficient than a mere Monte Carlo approach if the regularity of ξ compensates for the slowness of filling $\mathbb{X}$, which of course increases as the dimension d of $\mathbb{X}$ increases. By choosing the x_i s on a lattice, the fill distance can be made such that $h_n = O(n^{-1/d})$. Then, the convergence of $|A_u(\xi_n)|$ to $|A_u(\xi)|$ when the x_i s fill $\mathbb{X}$ regularly, is faster than Monte Carlo if $\nu > 3d/2$.

3 Convergence acceleration

3.1 Control of convergence

Of course, sampling $\mathbb{X}$ regularly as above may be suboptimal when the evaluations of f are sequential. This section addresses the problem of choosing a sequence $(x_n)_{n \in \mathbb{N}}$ so that the error of volume approximation conditioned on $\xi(x_i) = f(x_i)$, $i = 1, 2, \dots$ decreases rapidly. More precisely, a desirable strategy would consist in choosing

$$x_n = \operatorname{argmin}_{x_n \in \mathbb{X}} \Upsilon_n(x_n) := \mathbb{E} \left[(|A_u(\xi)| - |A_u(\xi_n)|)^2 \mid Z_{n-1} \right], \quad (16)$$

where for all n , $Z_n = (\xi(x_1), \dots, \xi(x_n))$. Note that $\Upsilon_n(x_n)$ can also be written as

$$\Upsilon_n(x_n) = \mathbb{E} \left[\mathbb{E} \left[(|A_u(\xi)| - |A_u(\xi_n)|)^2 \mid Z_n \right] \mid Z_{n-1} \right]. \quad (17)$$

The distribution of $|A_u(\xi)|$ conditioned on observations is generally unknown (see Adler, 2000, Section 4.4) and therefore, $\mathbb{E} \left[(|A_u(\xi)| - |A_u(\xi_n)|)^2 \mid Z_n \right]$ cannot be easily determined analytically. To overcome this difficulty, we could minimize a Monte Carlo approximation of (16) instead, namely

$$x_n = \underset{x_n \in \mathbb{X}}{\operatorname{argmin}} \Upsilon_{n,m}(x_n) := \mathbb{E} \left[m^{-1} \sum_{i=1}^m (|A_u(\xi_n + \zeta_n^i)| - |A_u(\xi_n)|)^2 \mid Z_{n-1}, \{\zeta_n^i, i \leq m\} \right], \quad (18)$$

where the random processes ζ_n^i are m independent copies of ξ conditioned on $Z_n = (0, \dots, 0)$. The program (18) becomes numerically tractable if we also replace $|A_u(\cdot)|$ by its Monte Carlo estimator $|A_u(\cdot)|_l$. Whereas simulating the conditioned processes ζ_n^i is easy in principle (see Chilès and Delfiner, 1999, chap. 7), it is also computationally intensive since it typically requires $O(l^3)$ operations to simulate ξ at given points $x_1, \dots, x_l$. Since l has to be high enough to ensure a degree of accuracy of the estimator $|A_u(\cdot)|_l$, conditional simulations ought to be avoided.

An alternative solution is to approximate $\mathbb{E} \left[(|A_u(\xi)| - |A_u(\xi_n)|)^2 \mid Z_n \right]$ by $\mathbb{E} \left[(|A_u(\xi)|_l - |A_u(\xi_n)|_l)^2 \mid Z_n, \{X_i, i \leq l\} \right]$, for l high enough. Then, the Minkowski inequality gives

$$\begin{aligned} & \mathbb{E} \left[(|A_u(\xi)|_l - |A_u(\xi_n)|_l)^2 \mid Z_n, \{X_i, i \leq l\} \right]^{1/2} \\ & \leq \frac{1}{l} \sum_{i=1}^l \mathbb{E} \left[(\mathbf{1}_{\xi(X_i) > u} - \mathbf{1}_{\xi_n(X_i) > u})^2 \mid Z_n, \{X_i, i \leq l\} \right]^{1/2}. \end{aligned} \quad (19)$$

This makes possible to build a *stepwise uncertainty reduction* algorithm as presented in the next section.

3.2 A stepwise uncertainty reduction algorithm

Denote by $S = \{y_1, \dots, y_l\}$ a set of l independent sample values of X . Given a finite sequence $(x_i)_{1 \leq i \leq n-1}$ of evaluation points, we wish to obtain a new point x_n that yields the largest decrease of the upper bound of the volume approximation mean-square error obtained in (19), i.e.,

$$x_n = \underset{x_n \in S}{\operatorname{argmin}} \Upsilon'_n(x_n) := \frac{1}{l} \sum_{i=1}^l \mathbb{E} \left[(\mathbf{1}_{\xi(y_i) > u} - \mathbf{1}_{\xi_n(y_i) > u})^2 \mid B_{n-1} \right]^{1/2}, \quad (20)$$

where B_n denotes the event $\{\xi(x_1) = f(x_1), \dots, \xi(x_n) = f(x_n)\}$, $n > 0$.

A few steps are needed to transform (20) into a numerically tractable program. First, note that

$$\begin{aligned} & \mathbb{E} \left[(\mathbf{1}_{\xi(y_i) > u} - \mathbf{1}_{\xi_n(y_i) > u})^2 \mid B_{n-1} \right] \\ & = \int_{z \in \mathbb{R}} \mathbb{E} \left[(\mathbf{1}_{\xi(y_i) > u} - \mathbf{1}_{\xi_n(y_i) > u})^2 \mid \xi(x_n) = z, B_{n-1} \right] \\ & \times p_{\xi(x_n) \mid B_{n-1}}(z) dz, \quad \forall i \in \{1, \dots, l\}, \end{aligned} \quad (21)$$

where $p_{\xi(x)|B_{n-1}}$ denotes the density of $\xi(x)$ conditionally to B_{n-1} . However, intrinsic Kriging assumes that the mean of ξ is unknown and therefore, for $x \in \mathbb{X}$, $\mathbb{E}[(\mathbf{1}_{\xi(x)>u} - \mathbf{1}_{\xi_n(x)>u})^2 | \xi(x_n) = z, B_{n-1}]$ cannot be determined exactly. Indeed, the values of $a_n(x)$, $b_n(x)$ and $\tilde{\sigma}_n(x)$ in (10) are unknown in practice. Nevertheless, (8) leads to the approximation

$$\mathbb{E}[(\mathbf{1}_{\xi(x)>u} - \mathbf{1}_{\xi_n(x)>u})^2 | \xi_n(x)] \approx v_n(x) := \Psi\left(\left|\frac{u - \xi_n(x)}{\sigma_n(x)}\right|\right). \quad (22)$$

Finally, define a discretization operator Δ_Q , which can be written for instance as

$$\forall h \in \mathbb{R}, \quad \Delta_Q h = z_1 + \sum_{i=2}^Q (z_i - z_{i-1}) \mathbf{1}_{[z_i, +\infty[}(h)$$

with $z_1 < z_2 < \dots < z_Q$. We can now write (20) as a numerically tractable program:

$$x_n = \underset{x_n \in S}{\operatorname{argmin}} \Upsilon_n''(x_n) := \frac{1}{l} \sum_{i=1}^l \left(\sum_{j=1}^Q \mathbb{P}\{\Delta_Q \xi(x_n) = z_j | B_{n-1}\} \mathbb{E}[v_n(y_i) | \xi(x_n) = z_j, B_{n-1}] \right)^{1/2}. \quad (23)$$

An informal interpretation of (23) is that x_n minimizes the error of prediction of $\mathbf{1}_{\xi(x)>u}$ by $\mathbf{1}_{\xi_n(x)>u}$, which is measured via $v_n(x)$, averaged on $\mathbb{X}$ under the distribution μ , and conditioned on the observations. When $\Upsilon_n''(x)$ becomes small for all $x \in S$, $|A_u(\xi_n)|_l$ conditioned on observations provides a good approximation of $\mathcal{P}_u$. As will be seen in Section 4, the proposed strategy is likely to achieve very efficient convergences.

4 Example

This section provides a one-dimensional illustration of the proposed algorithm. We wish to estimate (1), where $f(x)$ is a given function defined over $\mathbb{R}$ and $X \sim \mu = \mathcal{N}(0, \sigma^2)$. We assume that f is a sample path of ξ . After a few iterations, the unknown function f (as shown in Figure 1) has been sampled so that the probability of excursion $\mathbb{P}\{\xi(x) > u | \xi(x_i) = f(x_i), i = 1 \dots, n\}$ is determined accurately in the region where the probability density of X is high. This example illustrates the effectiveness of the proposed algorithm. Note that in practice, a parametrized covariance has to be chosen for ξ and its parameters should be estimated from the data, using, for instance, a maximum likelihood approach (e.g. Stein, 1999).

5 Appendix : Intrinsic Random Functions

In this section, we intend to summarize the most important notions about intrinsic random functions (Matheron, 1973). Let $\mathcal{N}$ be a vector space of functions $\{b^\top r(x), b \in \mathbb{R}^l\}$ and $\xi(x)$ be a random process with mean $m(x) \in \mathcal{N}$. The main idea of intrinsic random functions is to find some linear transformations of $\xi(x)$ *filtering out* the mean so as to consider a zero-mean process again.




Figure 1. Top: threshold u (horizontal solid line), function f (thin line), $n=10$ evaluations as obtained by the proposed algorithm using $l = 800$ and $Q = 20$ (squares), IK approximation f_n (thick line), 95% confidence intervals computed from the IK variance (dashed lines). Middle: probability of excursion (solid line), probability density of X (dotted line). Bottom: graph of $\Upsilon_n''(y_i)$, $i = 1, \dots, l = 800$, the minimum of which indicates where the next evaluation of f should be done (i.e., at approximately 0.75).

Let $\tilde{\Lambda}$ be the vector space of *finite-support measures*, i.e. the space of linear combinations $\sum_{i=1}^n \lambda_i \delta_{x_i}$, where δ_x stands for the Dirac measure, such that for any $B \subset \mathbb{X}$, $\delta_x(B)$ equals one if $x \in B$ and zero otherwise. Let $\tilde{\Lambda}_{\mathcal{N}^\perp}$ be the subset of the elements of $\tilde{\Lambda}$ that vanish on $\mathcal{N}$. Thus, $\lambda \in \tilde{\Lambda}_{\mathcal{N}^\perp}$ implies

$$\langle \lambda, f \rangle := \sum_{i=1}^n \lambda_i f(x_i) = 0, \quad \forall f \in \mathcal{N}.$$

In the following, we shall restrict ourselves to the case where $\mathcal{N}$ is a vector space of polynomials of degree at most equal to l . Denote by $\mathcal{N}_l$ the linear hull of all multivariate monomials x^i , where $i = (i_1, \dots, i_d)$ are multi-indexes such that $|i| := i_1 + \dots + i_d \leq l$, and define $\tilde{\Lambda}_l := \tilde{\Lambda}_{\mathcal{N}_l^\perp}$.

Let $\xi_G(\lambda)$ be a linear map on $\tilde{\Lambda}_l$, with values in $L^2(\Omega, \mathcal{A}, \mathbb{P})$, the space of second-order random variables. Assume that $\mathbb{E}[\xi_G(\lambda)] = 0$ for all λ and that

$$k(\lambda, \mu) := \text{cov}[\xi_G(\lambda), \xi_G(\mu)] = \sum_{i,j} \lambda_i \mu_j k(x_i, y_j),$$

where $k(x, y)$ is a symmetric conditionally positive definite function (i.e. a function such that $k(x, y) = k(y, x)$ and $k(\lambda, \lambda) \geq 0$ for all $\lambda \in \tilde{\Lambda}_l$). Then, $\xi_G(\lambda)$ is a *generalized random process* and $k(x, y)$ is called a *generalized covariance* (note that any covariance is a generalized covariance). Let $\tilde{\mathcal{H}}_l$ be the subspace of $L^2(\Omega, \mathcal{A}, \mathbb{P})$ spanned by $\xi_G(\lambda)$, $\lambda \in \tilde{\Lambda}_l$. Since random variables in $\tilde{\mathcal{H}}_l$ are zero-mean, the inner product of $L^2(\Omega, \mathcal{A}, \mathbb{P})$ can be expressed in $\tilde{\mathcal{H}}_l$ as

$$(\xi_G(\lambda), \xi_G(\mu))_{L^2(\Omega, \mathcal{A}, \mathbb{P})} = k(\lambda, \mu), \quad \lambda, \mu \in \tilde{\Lambda}_l.$$

Thus, the bilinear form $k(\lambda, \mu)$ endows $\tilde{\Lambda}_l$ and $\tilde{\mathcal{H}}_{\mathcal{N}^\perp}$ with a structure of pre-Hilbert space. The completions $\mathcal{H}_l$ and Λ_l of $\tilde{\mathcal{H}}_l$ and $\tilde{\Lambda}_l$ under this inner product define isomorphic Hilbert spaces. $\xi_G(\lambda)$ can be extended on Λ_l by continuity. Simplifying hypotheses are introduced in the next paragraph.

Let $\tau_h : \tilde{\Lambda}_l \rightarrow \tilde{\Lambda}$ be the translation operator such that for $\lambda = \sum_i \lambda_i \delta_{x_i} \in \tilde{\Lambda}_l$, $\tau_h \lambda = \sum_i \lambda_i \delta_{x_i+h}$. Note that $\tilde{\Lambda}_l$ is stable under translation since $\mathcal{N}_l$ is itself a translation-stable space of functions. Assume further that the generalized covariance $k(x, y)$ is invariant by translation. In the following, we shall write $k(h)$ with $h = x - y$ instead of $k(x, y)$, when the covariance is assumed to be stationary. Then τ_h is continuous and can be uniquely extended on Λ_l .

Definition 1. Let $\xi_G(\lambda)$ be a zero-mean generalized random process defined on Λ_l , with stationary generalized covariance $k(h)$. The random process $h \mapsto \xi_G(\tau_h \lambda)$, $\lambda \in \Lambda_l$, is therefore weakly stationary. $\xi_G(\lambda)$, $\lambda \in \Lambda_l$, is then an Intrinsic Random Function of order l , or *IRF(l)* in short.

If $\xi(x)$, $x \in \mathbb{X}$, is a second-order random process, with mean in $\mathcal{N}_l$ and covariance $k(x, y)$, the linear map

$$\begin{aligned} \xi : \tilde{\Lambda}_l &\rightarrow \mathcal{H} \\ \lambda = \sum_{i=1}^n \lambda_i \delta_{x_i} &\mapsto \xi(\lambda) := \sum_{i=1}^n \lambda_i \xi(x_i), \end{aligned}$$

extends $\xi(x)$ on $\tilde{\Lambda}_l$, where $\mathcal{H}$ stands for the Hilbert space generated by $\xi(x)$, $x \in \mathbb{X}$. Since $k(x, y)$ is positive definite, $(\lambda, \mu)_{\tilde{\Lambda}_l} := (\xi(\lambda), \xi(\mu))_{\mathcal{H}}$ defines an

inner product on $\tilde{\Lambda}_l$. Let Λ_l be the completion of $\tilde{\Lambda}_l$ under this inner product and extend $\xi(\lambda)$ on Λ_l by continuity (a generalized random process is thus obtained).

Definition 2. Let $\xi_G(\lambda)$ be an $IRF(l)$. A second-order random process $\xi(x)$, $x \in \mathbb{X}$, is a representation of $\xi_G(\lambda)$ iff

$$\xi_G(\lambda) = \xi(\lambda), \quad \forall \lambda \in \Lambda_l.$$

If $\xi_0(x)$ is any representation of $\xi_G(\lambda)$, other representations of $\xi_G(\lambda)$ can be written as

$$\xi(x) = \xi_0(x) + \sum_{i=1}^q B_i p_i(x), \quad (24)$$

where the p_i s form a basis of $\mathcal{N}_l$ and the B_i s are any second-order random variables. Thus, the representations of an $IRF(l)$ constitute a class of random processes with mean in $\mathcal{N}_l$ (Matheron, 1973).

References

- R. J. Adler. On excursion sets, tube formulas and maxima of random fields. *Ann. Appl. Probab.*, 10(1):1–74, 2000.
- J.-P. Chilès and P. Delfiner. *Geostatistics: Modeling Spatial Uncertainty*. Wiley, New York, 1999.
- P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events for Insurance and Finance*. Applications of Mathematics. Springer-Verlag, Berlin, 1997.
- G. S. Kimeldorf and G. Wahba. A correspondance between Bayesian estimation on stochastic processes and smoothing by splines. *Ann. Math. Statist.*, 41(2): 495–502, 1970.
- W. Light and H. Wayne. On power functions and error estimates for radial basis functions interpolation. *J. Approx. Theory*, 92(2):245–266, 1998.
- G. Matheron. The intrinsic random functions, and their applications. *Adv. Appl. Probab.*, 5:439–468, 1973.
- F. J. Narcowich, J. D. Ward, and H. Wendland. Refined error estimates for radial basis function interpolation. *Constr. Approx.*, 19(4):541–564, 2003.
- R. Y. Rubinstein. The cross-entropy method for combinatorial and continuous optimization. *Methodol. Comput. Appl. Probab.*, 2:127–190, 1999.
- M. L. Stein. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer, New York, 1999.
- H. Wendland. *Scattered Data Approximation*. Monographs on Applied and Computational Mathematics. Cambridge Univ. Press, Cambridge, 2005.
- Z. Wu and R. Schaback. Local error estimates for radial basis function interpolation of scattered data. *IMA J. Numer. Anal.*, 13:13–27, 1993.